


Product Name	Recombinant Bacteriophage T7 RNA Polymerase	<p>SDS-PAGE gel kDa 1 2</p> <p>235 170 130 93 70 53 41 30 22 18</p> <p>1 – MW Marker 2 – T7 RNA Pol</p>
Synonym(s)	DNA-directed RNA Polymerase	
Quantity	5,000 units, 25,000 units, 100,000 units	
Catalog Number	777627	
Concentration	50,000 Units/ml	
Molecular weight	100 kDa	
Purity	>95% by SDS-PAGE	
Tag	N-terminal 6xHis-tag	
Expression Source	E. coli	
GenBank Accession #	M38308.1, a. a. 2-882	
Application	RNA production for probe generation, vaccine development, RNA structural and/or catalytic investigation	
Formulation	50 mM Tris Cl pH 7.9, 130 mM NaCl, 20 mM BME, 1 mM EDTA, 0.1% Tween-20, 50% glycerol	
Storage and Stability	Stable for 12 months at -20°C, Avoid freeze/thaw cycles	
Description	Bacteriophage T7 RNA Polymerase is a 99 kDa protein that recognizes T7 phage promoters with high specificity and subsequently initiates transcription. T7 RNA polymerase is a single subunit, highly processive and stable enzyme, characteristics that make it suitable for a broad range of biochemistry and molecular biology applications.	
Reference	<ul style="list-style-type: none"> • Cold Spring Harb Protoc; 2013; doi:10.1101/pdb. prot078527 • Kartje ZJ, Janis HI, Mukhopadhyay S, Gagnon KT. J Biol Chem. 2021 Jan-Jun;296:100175. doi: 10.1074/jbc.RA120.014553. 	

T7 RNA Polymerase Activity



Components and Storage

Components	5000 U	25000 U	100000 U
T7 RNA Polymerase (50 U/ μ L)	100 μ L	500 μ L	2 mL
10X Reaction Buffer	100 μ L	500 μ L	2 mL

Store all the kit components at -20°C.

Protocol for Standard RNA Synthesis

Assemble the reaction at room temperature in the following order.

Components	Amount	Concentration
Nuclease-free water	X μ l	
10X Reaction Buffer	2 μ l	
NTP	X μ l	0.5 mM each
Template DNA	X μ l	0.2–1 μ g
RNase Inhibitor (optional)	0.5 μ l	1 U/ μ l final
Fresh DTT (optional)	X μ l	5 mM final

Components	Amount	Concentration
T7 RNA Pol	2 μ l	
Total Volume	20 μ l	

- 1) Mix thoroughly. Incubate at 37°C for 1 hour. For shorter (< 300 nt) transcripts incubate at 37°C for 2–16 hours.
- 2) (Optional) Add 1 μ L of DNase I to the reaction system and incubate at 37°C for 15min to digest the DNA template. Compared with the product RNA, the content of template DNA is very low. Generally, it does not need to be removed, and it can also be digested with DNase I.
- 3) Continue to purify the synthesized RNA or detect the transcription product by gel electrophoresis.

Products are for research use only and are not intended for human use. We do not sell to patients.